



PROTECTION FOR BOTH ADOLESCENTS AND ADULTS

Vaccinate against Pertussis, Tetanus, and Diphtheria

The number of Pertussis cases reported in 2004 was at a 45-year high. In fact, 25,827 cases of Pertussis were reported to the Centers for Disease Control and Prevention (CDC) that year.²

ADACEL vaccine is the first tetanus, diphtheria, and Acellular Pertussis (TDaP) booster vaccine available for both adolescents and adults 11 through 64 years of age.



SAFETY INFORMATION

ADACEL vaccine is indicated for active booster immunization for the prevention of tetanus, diphtheria, and Pertussis as a single dose in people 11 through 64 years of age. As with any vaccine, ADACEL vaccine may not protect 100% of people receiving the vaccine. There are risks associated with all vaccines. The most common local side effects with ADACEL vaccine were injection site pain, redness, and swelling. The most common systemic side effects were headache, body ache, tiredness, and fever. As with any vaccine, there is a small risk of allergic reactions. Some signs of allergic reactions are hives, swelling of the throat, low blood pressure, shock, and difficulty breathing. If you begin to experience any of these signs seek treatment right away. These reactions are rare and usually occur before leaving the doctor's office. When administering any intramuscular injection, like ADACEL vaccine, to people with bleeding disorders, caution should be exercised because they may develop a serious bruise or collection of blood at the injection site. If you notice any other problems or symptoms following vaccination, please contact your health-care professional immediately.

For more information about ADACEL vaccine, refer to the [Patient Information](#) on ADACELVACCINE.com